

DIFFERENTIAL SYSTEMS WITH DELAYS

SANDRA PINELAS

TUNISIA, 2022

A differential system

Let us to considerer the differential system

$$x^{(n)}(t) = \sum_{k=1}^n A_k x^{(n-k)}(t - \tau_k), \quad t \geq 0$$

where:

↪ n is even

↪ A_k are p -by- p real matrices

↪ τ_k are positive real numbers, for $k = 1, \dots, n$

The characteristic equation is

$$\lambda^n = \sum_{k=1}^n A_k \lambda^{n-k} e^{-\tau_k}$$

A differential equation

It is possible to prove that for

$$M(\lambda) = \sum_{k=1}^n A_k \lambda^{n-k} e^{-\tau_k}$$

The equation





$$x^{(n)}(t) = \sum_{k=1}^n A_k x^{(n-k)}(t - \tau_k), \quad t \geq 0$$

is oscillatory if and only if


$$\lambda^n \notin [-\mu(-M(\lambda)), \mu(-M(\lambda))]$$

A differential equation


$$\mu(M(\lambda)) < \lambda^n$$



A differential equation



$$\mu(-M(\lambda)) < -\lambda^n$$



$$\mu(M(\lambda)) < \lambda^n$$

Oscillatory system

Theorem 1: Let n even. If

$$\Rightarrow \mu(A_k) < 0$$

$$\Rightarrow \max_{\lambda < 0} \left\{ \lambda^{-n} e^{-\lambda \tau} \sum_{\substack{k=2 \\ k \text{ is even}}}^n \mu(A_k) - \lambda^{-1} e^{-\lambda T} \sum_{\substack{k=1 \\ k \text{ is odd}}}^{n-1} \mu(-A_k) \right\} < 1$$

then

$$x^{(n)}(t) = \sum_{k=1}^n A_k x^{(n-k)}(t - \tau_k), \quad t \geq 0$$

is oscillatory.


Oscillatory system

Draft of the proof:

$$\lambda = 0 \quad \longrightarrow \quad \mu(M(0)) = \mu(A_n) < 0$$

$$\lambda > 0 \quad \longrightarrow \quad \mu(M(\lambda)) \leq \sum_{k=1}^n \mu(A_k) \lambda^{n-k} e^{-\lambda \tau_k} < 0 < \lambda^n$$

$$\lambda < 0 \quad \longrightarrow \quad \frac{\mu(M(\lambda))}{\lambda^n} < 1 \quad \searrow$$

$$\frac{\mu(M(\lambda))}{\lambda^n} \leq \underbrace{\lambda^{-n} e^{-\lambda \tau} \sum_{\substack{k=2 \\ k \text{ is even}}}^n \mu(A_k)}_{f(\lambda)} - \lambda^{-1} e^{-\lambda T} \sum_{\substack{k=1 \\ k \text{ is odd}}}^{n-1} \mu(-A_k)$$


$$\lim_{\lambda \rightarrow 0^-} f(\lambda) = -\infty = \lim_{\lambda \rightarrow -\infty} f(\lambda)$$

Oscillatory system

Example 1: $x^{(4)}(t) = A_1 x^{(3)}(t - 0, 1) + A_2 x''(t - \tau_2) + A_3 x'(t - \tau_3) + A_4 x(t - 0, 4)$

⇒ $0, 1 < \tau_2, \tau_3 < 0, 4$

⇒ $A_1 = \begin{bmatrix} -8 & 1 \\ 4 & -10 \end{bmatrix} \quad \mu_1(A_1) = \max\{-4, -9\} = -4$

⇒ $A_2 = \begin{bmatrix} -7 & -3 \\ 2 & -4 \end{bmatrix} \quad \mu_1(A_2) = \max\{-5, -1\} = -1$

⇒ $A_3 = \begin{bmatrix} -3 & 4 \\ 0 & -10 \end{bmatrix} \quad \mu_1(A_3) = \max\{-3, -6\} = -3$

⇒ $A_4 = \begin{bmatrix} -5 & 3 \\ 4 & -6 \end{bmatrix} \quad \mu_1(A_4) = \max\{-1, -3\} = -1$

$$-2\lambda^{-4}e^{-0,1\lambda} + 7\lambda^{-1}e^{-0,4\lambda} < -7,673344$$

**The system is
oscillatory**

Oscillatory system

Theorem 2: Let n even. If

↳ for k even $\mu(A_k) < 0$

↳ for k odd $\mu(-A_k) < 0$

$$\text{↳ } \max_{\lambda > 0} \left\{ \lambda^{-n} e^{-\lambda T} \sum_{\substack{k=1 \\ k \text{ is even}}}^n \mu(A_k) - \lambda^{-1} e^{-\lambda \tau} \sum_{\substack{k=1 \\ k \text{ is odd}}}^n \mu(A_k) \right\} < 1$$

then

$$x^{(n)}(t) = \sum_{k=1}^n A_k x^{(n-k)}(t - \tau_k), \quad t \geq 0$$

is oscillatory.

Oscillatory system

Draft of the proof:

$$\lambda = 0 \quad \longrightarrow \quad \mu(M(0)) = \mu(A_n) < 0$$


$$\begin{aligned} \lambda < 0 \quad \longrightarrow \quad \mu(M(\lambda)) &\leq \sum_{\substack{k=2 \\ k \text{ is even}}}^n \mu(A_k) \lambda^{n-k} e^{-\lambda \tau_k} \\ &\quad - \sum_{\substack{k=1 \\ k \text{ is odd}}}^{n-1} \mu(-A_k) \lambda^{n-k} e^{-\lambda \tau_k} < 0 < \lambda^n \end{aligned}$$

Oscillatory system

Draft of the proof (continuation):

$$\lambda > 0 \quad \longrightarrow \quad \frac{\mu(M(\lambda))}{\lambda^n} < 1 \quad \searrow$$

$$\frac{\mu(M(\lambda))}{\lambda^n} \leq \underbrace{\lambda^{-n} e^{-\lambda T} \sum_{\substack{k=2 \\ k \text{ is even}}}^n \mu(A_k)}_{g(\lambda)} + \underbrace{\lambda^{-1} e^{-\lambda \tau} \sum_{\substack{k=1 \\ k \text{ is odd}}}^{n-1} \mu(A_k)}_{g(\lambda)}$$



$g(\lambda)$

$$\lim_{\lambda \rightarrow 0^+} g(\lambda) = -\infty$$

$$\lim_{\lambda \rightarrow +\infty} g(\lambda) = 0$$

Oscillatory system

Example 2: $x^{(4)}(t) = A_1 x^{(3)}(t-1) + A_2 x''(t-\tau_2) + A_3 x'(t-\tau_3) + A_4 x(t-0,25)$

⇒ $0,25 < \tau_2, \tau_3 < 1$

⇒ $A_1 = \begin{bmatrix} 1 & 0 \\ 0 & 1/2 \end{bmatrix} \quad \mu_\infty(A_1) = 1, \quad \mu_\infty(-A_1) = -1/2.$

⇒ $A_2 = \begin{bmatrix} -7 & -3 \\ 2 & -8 \end{bmatrix} \quad \mu_\infty(A_2) = -4.$

⇒ $A_3 = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} \quad \mu_\infty(A_3) = 1, \quad \mu_\infty(-A_3) = -1/12$

⇒ $A_4 = \begin{bmatrix} -6 & 4 \\ 1 & -9 \end{bmatrix} \quad \mu_\infty(A_4) = -2$

$$-6\lambda^{-4}e^{-\lambda} + 2\lambda^{-1}e^{-\lambda/4} < 0,655185324$$

**The system is
oscillatory**

Nonoscillatory system

$$\mu(-C) \leq 0 \Rightarrow \det(C) \geq 0;$$

if d is odd, $\mu(C) \leq 0 \Rightarrow \det(C) \leq 0$




$$x^{(n)}(t) = \sum_{k=1}^n A_k x^{(n-k)}(t - \tau_k)$$

is nonoscillatory if and only if there exist a λ_1 and λ_2 such that

$$\mu(-M(\lambda_1)) \leq -\lambda_1^n \quad \text{and} \quad \mu(M(\lambda_2)) \leq \lambda_2^n$$

Nonoscillatory system



Nonoscillatory system

Theorem 4: Let n even. If

↪ $\mu(-A_k) < 0$ for $k = 1, \dots, n$


then

$$x^{(n)}(t) = \sum_{k=1}^n A_k x^{(n-k)}(t - \tau_k)$$


is nonoscillatory independently of the delays.

Nonoscillatory system

Draft of the proof:

$\Rightarrow \lambda < 0$ 

$$\mu(-M(\lambda)) \leq \lambda^n \underbrace{\left(\frac{e^{-\lambda\tau}}{(-\lambda)^n} \sum_{\substack{k=1 \\ k \text{ even}}}^n \mu(-A_k) + \frac{e^{-\lambda T}}{-\lambda} \sum_{\substack{k=1 \\ k \text{ odd}}}^n \mu(A_k) \right)}_{f(\lambda)}$$



$$\lim_{\lambda \rightarrow 0^-} f(\lambda) = -\infty$$


$$f(\lambda) < -1 \implies \mu(-M(\lambda_1)) \leq -\lambda_1^n$$

There exist a $\lambda_1 < 0$ such that

Nonoscillatory system

Draft of the proof:

$$\Rightarrow \lambda > 0 \quad \longrightarrow \quad \mu(M(\lambda)) \leq \underbrace{\lambda^n \sum_{k=1}^n \mu(A_k) \lambda^{-k} e^{-\lambda \tau_k}}_{h(\lambda)}$$



$$\lim_{\lambda \rightarrow +\infty} h(\lambda) = 0^+$$

There exist a $\lambda_2 < 0$ such that

$$h(\lambda) < 1 \quad \longrightarrow \quad \mu(M(\lambda_2)) \leq \lambda_2^n$$

Nonoscillatory system

Example 4:

$$x^{(4)}(t) = A_1 x^{(3)}(t - \tau_1) + A_2 x''(t - \tau_2) + A_3 x'(t - \tau_3) + A_4 x(t - \tau_4)$$

$$A_1 = \begin{bmatrix} 6 & 0 & 1 \\ 1 & 4 & -1 \\ 1 & -2 & 7 \end{bmatrix}$$

$$\mu_{\infty}(-A_1) = -2$$

$$A_2 = \begin{bmatrix} 10 & 3 & -2 \\ 0 & 5 & 1 \\ -3 & 1 & 8 \end{bmatrix}$$

$$\mu_{\infty}(-A_2) = -4$$

$$A_3 = \begin{bmatrix} 2 & 0 & -1 \\ 1 & 6 & 0 \\ 0 & 4 & 9 \end{bmatrix}$$

$$\mu_{\infty}(-A_3) = -5$$

$$A_4 = \begin{bmatrix} 8 & 3 & 2 \\ -1 & 4 & -1 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\mu_{\infty}(-A_4) = -1$$

**The system is
nonoscillatory**

Nonoscillatory system

Theorem 5: Let n even.

↪ for k even $\mu(-A_k) < 0$

↪ for k odd $\mu(A_k) < 0$


then

$$x^{(n)}(t) = \sum_{k=1}^n A_k x^{(n-k)}(t - \tau_k)$$


is nonoscillatory independently of the delays.

Nonoscillatory system

Draft of the proof:

$\Rightarrow \lambda < 0$ 

$$\mu(-M(\lambda)) \leq \lambda^n \underbrace{\left(\lambda^{-n} e^{-\lambda \tau} \sum_{\substack{k=2 \\ k \text{ even}}}^n \mu(-A_k) - \lambda^{-1} e^{-\lambda T} \sum_{\substack{k=1 \\ k \text{ odd}}}^{n-1} \mu(-A_k) \right)}_{f(\lambda)}$$




$$\lim_{\lambda \rightarrow 0^-} f(\lambda) = -\infty$$

There exist a $\lambda_1 < 0$ such that


$$f(\lambda) < -1 \implies \mu(-M(\lambda_1)) \leq -\lambda_1^n$$

Nonoscillatory system

Draft of the proof:

$\Rightarrow \lambda > 0$ 

$$\mu(M(\lambda)) \leq \lambda^n \underbrace{\left(\lambda^{-1} e^{-\lambda \tau} \sum_{\substack{k=2 \\ k \text{ even}}}^n \mu(A_k) + \lambda^{-n} e^{-\lambda T} \sum_{\substack{k=1 \\ k \text{ odd}}}^{n-1} \mu(A_k) \right)}_{h(\lambda)}$$



$$\lim_{\lambda \rightarrow +\infty} h(\lambda) = 0^+$$

There exist a $\lambda_2 < 0$ such that

$$h(\lambda) < 1 \quad \Rightarrow \quad \mu(M(\lambda_2)) \leq \lambda_2^n$$

Nonoscillatory system

Example 5:

$$x^{(4)}(t) = A_1 x^{(3)}(t - \tau_1) + A_2 x''(t - \tau_2) + A_3 x'(t - \tau_3) + A_4 x(t - \tau_4)$$

$$A_1 = \begin{bmatrix} -7 & 1 & -1 \\ 0 & -6 & -1 \\ 3 & 2 & -5 \end{bmatrix}$$

$$\mu_1(A_1) = -3$$

$$A_2 = \begin{bmatrix} 6 & 0 & 3 \\ 1 & -4 & 0 \\ 0 & 2 & 5 \end{bmatrix}$$

$$\mu_1(-A_2) = -2$$

$$A_3 = \begin{bmatrix} -9 & 0 & 1 \\ -1 & -1 & 0 \\ 3 & 0 & -2 \end{bmatrix}$$

$$\mu_1(A_3) = -1$$

$$A_4 = \begin{bmatrix} 10 & -3 & 0 \\ 2 & 5 & 3 \\ 0 & 1 & 6 \end{bmatrix}$$

$$\mu_1(-A_4) = -1$$

**The system is
nonoscillatory**

Nonoscillatory system

Theorem 6: Let n even

↳ for $k = 1, \dots, i$ $\mu(-A_k) < 0$

↳ for $k = i+1, \dots, n$ $\mu(A_k) < 0$

then

$$x^{(n)}(t) = \sum_{k=1}^n A_k x^{(n-k)}(t - \tau_k)$$

is nonoscillatory independently of the delays.

Nonoscillatory system

Draft of the proof:

$$\begin{aligned}
 \Rightarrow \lambda < 0 & \longrightarrow \mu(-M(\lambda)) \\
 & \leq \lambda^n \left[(-\lambda)^{-2} e^{-\lambda T^*} \sum_{\substack{k=2 \\ k \text{ is even}}}^i \mu(-A_k) + \lambda^{-n} e^{-\lambda \tau^*} \sum_{\substack{k=i+1 \\ k \text{ is even}}}^n \mu(-A_k) \right. \\
 & \quad \left. - \lambda^{-i} e^{-\lambda \tau} \sum_{\substack{k=1 \\ k \text{ is odd}}}^i \mu(A_k) - \lambda^{-(i+1)} e^{-\lambda T} \sum_{\substack{k=i+1 \\ k \text{ is odd}}}^{n-1} \mu(A_k) \right] \\
 & \quad \underbrace{\hspace{15em}}_{f(\lambda)} \\
 & \quad \lim_{\lambda \rightarrow 0^-} f(\lambda) = -\infty
 \end{aligned}$$

There exist a $\lambda_1 < 0$ such that $f(\lambda) < -1 \longrightarrow \mu(-M(\lambda_1)) \leq -\lambda_1^n$


Nonoscillatory system

Draft of the proof:

↪ $\lambda > 0$



$$\mu(M(\lambda)) \leq \lambda^n \left(\lambda^{-i} e^{-\lambda T} \sum_{k=1}^i \mu(A_k) + \lambda^{-(i+1)} e^{-\lambda \tau} \sum_{k=i+1}^n \mu(A_k) \right)$$



$h(\lambda)$

$$\lim_{\lambda \rightarrow +\infty} h(\lambda) = 0^+$$

There exist a $\lambda_2 < 0$ such that $h(\lambda) < 1 \implies \mu(M(\lambda_2)) \leq \lambda_2^n$

Nonoscillatory system

Example 6:

$$x^{(6)}(t) = \sum_{k=1}^5 A_k x^{(n-k)}(t - \tau_k)$$

**The system is
nonoscillatory**

$$A_1 = \begin{bmatrix} -5 & 1 & 0 \\ 3 & -6 & -4 \\ 0 & 2 & -9 \end{bmatrix} \quad A_2 = \begin{bmatrix} -8 & 1 & -2 \\ 1 & -6 & 3 \\ -1 & 0 & -10 \end{bmatrix} \quad \begin{aligned} \mu_1(A_1) &= -2 \\ \mu_1(A_2) &= -5 \end{aligned}$$

$$A_3 = \begin{bmatrix} -8 & 3 & 0 \\ 1 & -4 & 1 \\ 3 & 0 & -2 \end{bmatrix} \quad A_4 = \begin{bmatrix} -3 & 2 & -1 \\ 0 & -5 & 0 \\ 1 & 0 & -6 \end{bmatrix} \quad \begin{aligned} \mu_1(A_3) &= -1 \\ \mu_1(A_4) &= -2 \end{aligned}$$

$$A_5 = \begin{bmatrix} 6 & 4 & -1 \\ -1 & 7 & 0 \\ 2 & 0 & 8 \end{bmatrix} \quad A_6 = \begin{bmatrix} 5 & 0 & 1 \\ 0 & 4 & 2 \\ 1 & 2 & 6 \end{bmatrix} \quad \begin{aligned} \mu_1(-A_5) &= -3 \\ \mu_1(-A_6) &= -2 \end{aligned}$$

References

- ↪ I. Györi and G. Ladas, Oscillation Theory of Delay Differential Equations. Oxford Univ. Press, 1991.
- ↪ R. P. Agarwal, S.R. Grace and D. O'Reagan, Oscillation Theory for Difference and Functional Differential Equations. Kluwer, 2000.
- ↪ C.A. Desoer and M. Vidyasagar, Feedback Systems: Input-Output Properties. Ac. Press, 1975.
- ↪ José M. Ferreira e Sandra Pinelas, Oscillatory mixed difference systems, Advances in Difference Equations, vol. 2006, Article ID 92923 (2006), 1-18.
- ↪ J. Kirchner and U. Stroinsky, Explicit oscillation criteria for systems of neutral equations with distributed delay. Differential Equations and Dynam. Systems 3: 101-120 (1995).

References

- ✚ Sandra Pinelas, Nonoscillations in odd order difference systems of mixed type, Proceedings of the International Conference on Difference Equations, Special Functions and Applications, Munique 2005, S. Elaidy, J. Cushing, R. Lassr, V. Papageorgou, A. Rung, W. Van Assche (eds.), World Scientific, 2006, 507-519
- ✚ T. Krisztyn, Nonoscillations for functional differential equations of mixed type, Journal of Mathematical Analysis and Applications 254 (2000) 326-345